H-theorem in quantum systems interacting with reservoir

G. B. Lesovik,[1] A. V. Lebedev,[2] I. A. Sadovskyy,[3] M. V. Suslov,[4] and V. M. Vinokur[3]

[1] *L.D. Landau Institute for Theoretical Physics RAS, Akad. Semenova av., 1-A, Chernogolovka, 142432, Moscow Region, Russia*

[2] *Theoretische Physik, Wolfgang-Pauli-Strasse 27, ETH Zürich, CH-8093 Zürich, Switzerland*

[3] *Materials Science Division, Argonne National Laboratory, 9700 S. Cass Av., Argonne, IL 60637, USA*

[4] *Moscow Institute of Physics and Technology, Institutskii per. 9, Dolgoprudny, 141700, Moscow District, Russia*

(Dated: December 23, 2015)

Building on the concept of a quantum channel, a trace-preserving completely positive map that describes a temporary evolution of an open quantum system density matrix, we derive the criterion, which the system evolution operator should meet for the system to evolve with the non-diminishing entropy. This enables us to formulate quantum H-theorem in terms of observable quantities for a system interacting with the reservoir. We apply our criterion to generic physical quantum systems and uncover the conditions under which these systems follow quantum H-theorem. We discuss manifestation of the second law of thermodynamics in quantum physics and uncover special situations where it can be violated.

arXiv:1407.4437v2 [quant-ph] 22 Dec 2015

The foundational Boltzmann's H-theorem[1,2] states that the evolution of system described by the distribution function obeying the kinetic equation occurs with non-diminishing entropy. Having defined the entropy through the quantum mechanical density matrix $\hat{\rho}$ as $S(\hat{\rho}) = -k_{\rm B}\mathrm{tr}\{\hat{\rho}\log\hat{\rho}\}$, von Neumann conjectured[3] a solely quantum origin of the H-theorem. The proof of this conjecture became one of the major fundamental challenges.[4] The decades-long pursuit was crowned by recent progress of quantum information theory (QIT) that formulated several rigorous mathematical theorems about the conditions for a non-negative entropy gain.[5,6] Yet, the relation of these results to realm of physics remained not thoroughly understood. Here, building on the mathematical formalism provided by the QIT, we formulate the quantum H-theorem in terms of physical observables and discuss manifestation of the second law of thermodynamics in quantum physics. In what follows we uncover special situations where the second law can be violated and demonstrate that the typical evolution of energy-isolated quantum systems occurs with the non-diminishing entropy.

To describe quantum dynamics of an open system, the quantum information theory introduces the so-called *quantum channel* (QC) defined as a trace-preserving completely positive map, $\hat{\tilde{\rho}} = \Phi(\hat{\rho})$, of a density matrix.[5] The entropy gain in a channel $\Phi(\hat{\rho})$ is[7]

$$S(\Phi(\hat{\rho})) - S(\hat{\rho}) \geqslant -k_{\rm B}\mathrm{tr}\{\Phi(\hat{\rho})\log\Phi(\hat{1})\}, \qquad (1)$$

where $\hat{1}$ is the unit matrix. The most general class of the channels for which the entropy gain $\Delta S \equiv S(\hat{\tilde{\rho}}) - S(\hat{\rho}) \geqslant 0$, is the so-called *unital* channel defined by the relation[5,6] $\Phi(\hat{1}) = \hat{1}$, so that $\mathrm{tr}\{\Phi(\hat{\rho})\log\Phi(\hat{1})\} = 0$. Then within the framework of the QIT one can formulate the quantum H-theorem as follows: the entropy gain during evolution is non-negative if and only if the system evolution can be described by the unital channel.[?] To connect this formulation to the realm of physics note that any quantum system interacting with the reservoir generates a quantum channel. Indeed, let us consider joint evolution of the grand system, comprising a given quantum system and a reservoir initially prepared in a disentangled state,[?] $\hat{\mathcal{P}} = \hat{\rho}\otimes\hat{\rho}_{\rm R}$, where $\hat{\rho}_{\rm R}$ is the density matrix of the reservoir. Let $\hat{U}$ be the unitary operator describing the temporal evolution of the grand system. Then, according to the Stinespring-Kraus dilation theorem[5] $\Phi(\hat{\rho}) = \mathrm{tr}_{\rm R}\{\hat{U}\hat{\mathcal{P}}\hat{U}^{\dagger}\}$ is the quantum channel.

In physics the positive entropy gain, according to the second law of thermodynamics, is ensured by the energy isolation of the evolving system, see detailed discussion in Appx. A. Accordingly, in what follows we discuss the systems energy-isolated from the reservoir. Furthermore, we will be assuming that our systems are initially disentangled from the reservoir. Let us consider a fixed energy subspace E of the system Hilbert space spanned by the orthonormal basis states $|\psi_{i,E}\rangle$, where index i denotes all the remaining system's degrees of freedom. It is convenient to present the evolution operator, $\hat{U}$, of the grand system (a system plus reservoir) as

$$\hat{U} = \sum\nolimits_{E} |\psi_{j,E}\rangle\langle\psi_{i,E}|\, s_{ji,E}\hat{F}_{ji,E}, \qquad (2)$$

The factorization into $s_{ji,E}$ and $\hat{F}_{ji,E}$ is not unique, so we will be choosing the most suitable one for each particular case. For example, considering below a particle interacting with a spin, we will let $s_{ji,E}$ be the components of the scattering matrix corresponding to the transition amplitude between the particle states $|\psi_{i,E}\rangle \to |\psi_{j,E}\rangle$ (without taking into account interaction with the spin).

Operators $\hat{F}_{ji,E}$ are the family of operators acting in the reservoir Hilbert space, where subscripts i, j, and E specify the particle state. For an energy-isolated quantum system the quantum states at different energies transform independently. To determine whether the evolution belongs in the class of the unitality channel, one has to check if the system obeys the $\Phi(\hat{1}) = \hat{1}$ relation. Using the unitarity of $\hat{U}$, one finds (see Apps. A and B)

$$\Phi_{jj'}(\hat{1}_E) = [\hat{1}_E]_{jj'} + \sum\nolimits_{i} s_{ji,E}s^{*}_{j'i,E}\langle[\hat{F}^{\dagger}_{j'i}, \hat{F}_{ji}]\rangle, \qquad (3)$$

where $\langle \dots \rangle$ is averaging with respect to the initial state of the reservoir, and $\hat{1}_{E} = \sum_i |\psi_{j,E}\rangle\langle\psi_{i,E}|$. This relation is our central result. It establishes the criterion for unitality of the energy-isolated system in terms of physical operators describing the interaction of the quantum system with the reservoir and allows to formulate quantum H-theorem as follows:

Let the quantum system interacting with the reservoir be initially disentangled from it and be energy-isolated during the evolution. Let $\sum_i s_{ji,E} s^*_{j'i,E} \langle [\hat{F}^\dagger_{j'i}, \hat{F}_{ji}] \rangle = 0$, *where operators* $\hat{F}_{ji,E}$ *and coefficients* $s_{ji,E}$ *are defined as in Eq. (2), for energies* E *at which the system can be found with finite probability, i.e. for energies such that* $\langle \psi_{j,E}| \hat{\rho} |\psi_{j,E}\rangle \neq 0$. *Then the entropy gain* $\Delta S = S(\Phi(\hat{\rho})) - S(\hat{\rho}) \geqslant 0$.

The obtained formulation of the quantum H-theorem enables us to reveal a fundamental difference in how the second law of thermodynamics manifests itself in quantum and classical physics. In classical thermodynamics the energy-isolated system inevitably evolves with non-diminishing entropy. We find that in quantum physics it is different. To demonstrate that, we construct an energy-isolated quantum system for which $\langle [\hat{F}^\dagger_{j'i}, \hat{F}_{ji}] \rangle \neq 0$ and which thus evolves with the negative entropy gain. Let us consider a charged particle moving in a three-lead conductor and interacting with the spin via the induced magnetic field, see Fig. 1, and, according to our general framework, initially disentangled from the spin. In the absence of the external magnetic field the energy exchange is absent and the particle is energy-isolated. The joint scattering states of the particle and spin have the form

$$|\psi^{(\mathrm{in})}_\alpha\rangle|\sigma_0\rangle \to \sum\nolimits_\beta s_{\beta\alpha} |\psi^{(\mathrm{out})}_\beta\rangle \hat{U}_\beta \hat{U}^\dagger_\alpha |\sigma_0\rangle, \qquad (4)$$

where $|\psi^{(\mathrm{in/out})}_\alpha\rangle$ is the particle's incoming/outgoing state in the lead α, $s_{\beta\alpha}$ are the components of the scattering matrix of the three lead set up, $|\sigma_0\rangle$ is the initial state of the spin and $\hat{U}_\alpha$ ($\hat{U}^\dagger_\alpha$) is the unitary spin-1/2 rotation of the spin due to outgoing (incoming) electron in the lead α. Then the operators $\hat{F}$ of Eq. (3) are defined as $\hat{F}_{\beta\alpha} = \hat{U}_\beta \hat{U}^\dagger_\alpha$.

Let us recall now that rotations of a spin about different axis in general do not commute. We choose spin-1/2 rotations as $\hat{U}_1 = \hat{1}$, $\hat{U}_2 = i\hat{\sigma}_x$ and $\hat{U}_3 = i\hat{\sigma}_y$, where $\hat{\sigma}_x$ and $\hat{\sigma}_y$ are the Pauli matrices, so that $[\hat{U}_\alpha, \hat{U}_\beta] \neq 0$. Accordingly, $\langle [\hat{F}^\dagger_{\beta'\alpha}, \hat{F}_{\beta\alpha}] \rangle \neq 0$, and the resulting quantum channel is non-unital. The explicit calculation gives (see Appx. C)

$$\begin{aligned}\Phi(\hat{1}_{E}) = \hat{1}_{E} + 2i \big\{ & |1\rangle\langle 2| \, s_{13,E} s^*_{23,E} \langle \hat{\sigma}_x \rangle + \\ & |1\rangle\langle 3| \, s_{12,E} s^*_{23,E} \langle \hat{\sigma}_y \rangle - \\ & |2\rangle\langle 3| \, s_{12,E} s^*_{13,E} \langle \hat{\sigma}_z \rangle - \text{H.c.} \big\}. \end{aligned} \qquad (5)$$

Let the initial state of the spin be a pure state $|S_0\rangle = (|\uparrow_x\rangle + |\uparrow_y\rangle + |\uparrow_z\rangle)/\sqrt{3}$, so that all the $\langle \hat{\sigma}_\alpha \rangle$ in the Eq. (5) are equal to 1/3. Hence all off-diagonal elements of $\Phi(\hat{1}_{E})$ appear finite. Taking all $s = 2/3$, at some energy E_0, we construct the normalized initial state of the particle as $\hat{\rho} = (1/6) \int dE \, f(E - E_0) \hat{1}_{E}$, with $f(E)$ being the normalized to unity distribution function centered around $E = 0$ and rapidly decaying as $|E| \to \infty$, and obtain $\Delta S \approx -0.05 \, k_{\mathrm{B}}$. We thus demonstrated that even the energy isolation does not guarantee the evolution with the non-diminishing entropy. Note that in the discussed example the reservoir acts as some quantum analogue of the classical Maxwell demon. Namely, having been prepared in a special state, the reservoir is able to decrease the entropy of the system without the energy exchange with it, and can be referred to as a 'quantum Maxwell demon.' The applications of the quantum Maxwell demon concept for making quantum thermal machines will be published elsewhere.



Figure 1. Scattering in a 3-lead setup. A particle incident from the lead 1 is scattered into two other leads 2 and 3. Propagating particle induces magnetic field perpendicular to the lead direction. The spin is placed at the point where the respective fields induced by particles propagating along leads 2 and 3 are perpendicular to each other. To simplify consideration, we choose the set up design allowing to neglect the field induced by the particle in the lead 1.

As a first example of a system satisfying quantum H-theorem, we consider an electron elastically scattered by the one-dimensional (1D) potential barrier, see Figs. 2(a) and 2(b). Let the electron reflection be accompanied by the change of the reservoir state (e.g. let the reflection to imply emission of the low energy photon via braking radiation, or scattering of the photon), see Fig. 2(b) and the electron transmission to retain the state of the reservoir, see Fig. 2(a). Accordingly, the joint scattering states of the particle with the fixed energy assume the form,

$$|\mathrm{L}\rangle|n\rangle \to s_{\mathrm{RL}}|\mathrm{R}\rangle|n\rangle + s_{\mathrm{LL}}|\mathrm{L}\rangle|n'\rangle, \qquad (6)$$

$$|\mathrm{R}\rangle|n\rangle \to s_{\mathrm{LR}}|\mathrm{L}\rangle|n\rangle + s_{\mathrm{RR}}|\mathrm{R}\rangle|n''\rangle, \qquad (7)$$

where indices $\{\mathrm{L}, \mathrm{R}\}$ denote the incident (scattered) states in the left and right leads, respectively, $|n\rangle$ is the initial reservoir state, and $|n'\rangle$ and $|n''\rangle$ are the reservoir states resulting from the backscattering to the left and to the right, respectively. The unitality of the corresponding channel follows now from general Eq. (3), see Appx. D. However, for illustrative purpose we derive unitality straightforwardly using the explicit form of (6) and



Figure 2. Braking radiation in 1D. (a) A scattering where electron is transmitted without photon emission. (b) A backward scattering event accompanied by the emission of photons.

(7). Namely, calculating $\hat{1}_E$ in the basis of left and right scattering channels, $i, j \in \{\mathrm{L}, \mathrm{R}\}$, we obtain

$$\hat{1}_E \to |\mathrm{R}\rangle\langle\mathrm{R}|\,(|s_{\mathrm{RR}}|^2 + |s_{\mathrm{RL}}|^2) + |\mathrm{L}\rangle\langle\mathrm{L}|\,(|s_{\mathrm{LR}}|^2 + |s_{\mathrm{LL}}|^2) + \left\{|\mathrm{R}\rangle\langle\mathrm{L}|\,(s^*_{\mathrm{LR}} s_{\mathrm{RR}}\langle n''|n\rangle + s^*_{\mathrm{LL}} s_{\mathrm{RL}}\langle n|n''\rangle) + \mathrm{H.c.}\right\} \quad (8)$$

and taking into account the unitarity of the overall transformation, we arrive at $\Phi(\hat{1}_E) = \hat{1}_E$. Since this condition holds for any E, then for any state the system evolves with $\Delta S \geqslant 0$. The above consideration with some minor modifications holds for the grand system where the role of the reservoir is taken up by a single spin located near the scatterer. Then the spin remains intact if the particle is reflected and is rotated by the magnetic field induced by the transmitted particle.

Let us generalize the above consideration onto the particle propagating along the two-dimensional array of scatterers and spins (comprising the reservoir) located in the xy-plane. The magnetic field induced by the propagating particle is perpendicular to the plane, and all spins experience the commuting unitary rotations around the perpendicular z-axis. All the rotations commute, hence the condition of the quantum H-theorem is satisfied and $\Delta S \geqslant 0$.

Remarkably, the property of unitality of the grand system with the spin reservoir preserves in the three-dimensional (3D) case. As we mentioned above, the rotations experienced by an individual spin may appear, in general, non-commuting. Note, however, that operators $\hat{F}$ in Eq. (3) acquire the form

$$\hat{F}_{ji} = \hat{U}^{(1)}_{ji} \otimes \cdots \otimes \hat{U}^{(N)}_{ji}, \quad (9)$$

where $\hat{U}^{(a)}_{ji}$ is a unitary rotation of a spin a by the electron experienced a scattering process from the state $|i\rangle$ to the state $|j\rangle$. Then

$$[\hat{F}^\dagger_{j'i}, \hat{F}_{ji}] = \prod_{a=1}^{N} \otimes\, \hat{U}^{\dagger(a)}_{j'i}\hat{U}^{(a)}_{ji} - \prod_{a=1}^{N} \otimes\, \hat{U}^{(1)}_{ji}\hat{U}^{\dagger(1)}_{j'i}, \quad (10)$$

where $N \gg 1$ is the total number of spins. For most of the spins, the factors that appear upon averaging (10)



Figure 3. 1D random walk of an electron. Two-level systems (TLSs) shown as double well potentials are located equidistantly along the wire. Each TLS forms an effective potential for the electron, which depends on the TLS's quantum state. For simplicity we consider a completely transparent (open) or completely reflective (closed) effective scattering potential depending on the TLS state. At each scattering event the set of TLSs is replaced by a new (unentangled) one.

are small, $\langle \hat{U}^{\dagger(a)}_{j'i}\hat{U}^{(a)}_{ji}\rangle$, $\langle \hat{U}^{(a)}_{ji}\hat{U}^{\dagger(a)}_{j'i}\rangle < q < 1$. Therefore, we arrive at the estimate

$$\langle [\hat{F}^\dagger_{j'i}, \hat{F}_{ji}]\rangle \lesssim \exp[-N\ln(1/q)], \quad (11)$$

therefore, $\langle [\hat{F}^\dagger_{j'i}, \hat{F}_{ji}]\rangle \to 0$ as $N \to \infty$. We see that in the macroscopic limit of the number of spins, the averaged commutators appearing in the condition for the H-theorem vanish in spite of the fact that the commutators for the individual spins could remain finite. Thus the evolution of the 3D system involved occurs with $\Delta S \geqslant 0$.

To proceed further, we note, that if one can find a basis in the reservoir Hilbert space where all $\hat{F}_{ji}$-operators are diagonal, then the operators $\hat{F}_{ji}$ commute. Below we present two generic physical examples where this basis can be found explicitly: (i) an electron interacting with adiabatic two-level impurities and (ii) electron-phonon interaction in solids.

Consider the electron scattering on impurities that fluctuate between two positions with nearly equal energies. To be concrete we focus on the random walk of the electron along the ensemble of TLS (see the description of the similar systems in Refs. 12 and 13) as shown in Fig. 3. Assume for simplicity that (i) the dynamics of a TLS is slow hence its state does not change during the interaction with the electron and that (ii) each TLS which is in the state $|\uparrow\rangle$ ($|\downarrow\rangle$), scatters the electron via elastic unitary matrix $\hat{S}_\uparrow$ ($\hat{S}_\downarrow$). Then the global unitary transformation is given by $\hat{U} = \prod_n \otimes \hat{U}_n$ where $\hat{U}_n = \hat{S}^{(n)}_\uparrow |\uparrow_n\rangle\langle\uparrow_n| + \hat{S}^{(n)}_\downarrow |\downarrow_n\rangle\langle\downarrow_n|$ describes the scattering on the n-th impurity. Denote the scattering state of the electron moving in a direction $\mathbf{k}$ by $|\mathbf{k}\rangle$. Then the reservoir operator $\hat{F}_{\mathbf{k}\mathbf{k}'} = \prod_n \otimes \hat{F}^{(n)}_{\mathbf{k}\mathbf{k}'}$ with $\hat{F}^{(n)}_{\mathbf{k}\mathbf{k}'} = [\hat{S}^{(n)}_\uparrow]_{\mathbf{k}\mathbf{k}'}|\uparrow_n\rangle\langle\uparrow_n| + [\hat{S}^{(n)}_\downarrow]_{\mathbf{k}\mathbf{k}'}|\downarrow_n\rangle\langle\downarrow_n|$, see Eq. (2), commute with each other and, therefore, the second term in Eq. (3) vanishes. Hence, in each particular scattering event the entropy of the particle is non-diminishing.



Figure 4. Electrons and phonons in an atomic lattice. The scattering state of an electron depends strongly on the position of the scattering atom in the lattice. Importantly, the position of the scatterer remains almost unchanged during the scattering process because of the significant mass difference between the scattered electron and the scattering atom.

The question that now arises is whether the non-diminishing entropy maintains for the sequence of scatterings. Recall that in order for the entropy to grow monotonously, an electron should be disentangled from the TLS with which it is going to interact. Since in the course of the evolution an electron may return to the TLS on which it scattered and, therefore, got entangled in the past, these returns would violate this 'initial non-entanglement condition.' Thus to guarantee the evolution with non-diminishing entropy the TLS should have interacted with some other degrees of freedom that would lead to the memory loss of this TLS prior to the possible return of the electron. This memory loss is the manifestation of the so-called *monogamy* of entanglement[14] which is a specific property of the entanglement distribution between quantum systems: if a TLS is already entangled with an electron and later it becomes entangled with another degree of freedom then the initial entanglement with the electron vanishes. Thus the process of sequential scattering of an electron satisfies the H-theorem if the typical entanglement time for the TLS is less than the typical return time of an electron to particular TLS.

Now we demonstrate that the electron-phonon interaction leads to the electron evolution which satisfies the conditions of quantum H-theorem. In a standard electron-phonon interaction model, a representative test electron 'sees' a screened short-range ionic potential. Since the resulting scattering time is short and the ion is much heavier than the electron, the position of a given ion remains nearly unchanged during the interaction with the electron, see Fig. 4. Then the standard consideration of the electron-phonon interaction[15] results in a conclusion that at high temperatures where the typical energy of an electron is relatively high (i.e. it exceeds the Debye energy, the maximal energy of phonons), the electron-phonon collisions are quasi-elastic. This allows us to apply the same arguments as for the model of an electron interacting with two-level impurities considered above. Indeed, the slow ion-dynamics preserves the classical distribution function, $\rho_{\rm ion}(\{\mathbf{r}\})$, for the ions positions $\{\mathbf{r}\} = \{\mathbf{r}_1, \mathbf{r}_2, \ldots\}$ and hence the induced $\hat{F}$-operators are diagonal in the ion coordinate basis, $\hat{F}_{ji} = \int \prod_\alpha d\mathbf{r}_\alpha \, s_{ji}(\{\mathbf{r}\})\rho_{\rm ion}(\{\mathbf{r}\}) \prod_\alpha |\mathbf{r}_\alpha\rangle\langle\mathbf{r}_\alpha|$, where $s_{ji}(\{\mathbf{r}\})$ are electron scattering matrix on the potential defined by the ions positions $\{\mathbf{r}\}$. Therefore the conditions of the quantum H-theorem hold for electrons with energies exceeding the Debye energy. The initial disentanglement of the electron from the ionic reservoir is ensured either by the fact that a given ion did not participate in the previous collisions with the electron, or has already 'forgotten' about such an event due to the monogamy of entanglement.

It is noteworthy that unlike the other systems discussed in previous sections, the evolution of the electron-phonon system does not occur in the unital channel. Nevertheless, it satisfies the condition of the H-theorem thus illustrating a mathematical result establishing that for some classes of initial states even the evolution defined by non-unital channels may occur with the non-diminishing entropy.[9]

We are delighted to thank G. Blatter, G.-M. Graf, L. B. Ioffe and G. G. Amosov for illuminating discussions. The work was supported by the RFBR Grant No. 14-02-01287 (G. L.), by the Pauli Center for Theoretical Studies at ETH Zurich (G. L.), by the Swiss National Foundation through the NCCR QSIT (A. L.), and by the U.S. Department of Energy, Office of Science, Materials Sciences and Engineering Division (I. S. and V. V.).

Appendix A: Proof of the H-theorem

Here we derive Eq. (3) and sketch the proof of the H-theorem. We consider a quantum system which interacts with a quantum reservoir. The evolution operator $\hat{U}$ is unitary, $\hat{U}\hat{U}^\dagger = 1$, and using Eq. (2) we write the unitarity constraints in the form,

$$\sum\nolimits_i s_{ji,E} s^*_{j'i,E} \, \langle m|\hat{F}_{ji,E}\hat{F}^\dagger_{j'i,E}|m'\rangle = \delta_{jj'}\delta_{mm'}. \qquad \text{(A1)}$$

Let the reservoir initially be in a state described by the density matrix $\hat{\pi} = \sum_{nn'} \pi_{nn'} \, |n\rangle\langle n'|$. Then

$$[\Phi(\hat{\rho})]_{jj'} = \sum\nolimits_{ii'} \rho_{ii'} \mathrm{tr}\big\{ \langle\psi_j|\hat{U}|\psi_i\rangle \, \hat{\pi} \, \langle\psi_{i'}|\hat{U}^\dagger|\psi_{j'}\rangle \big\} \qquad \text{(A2)}$$

where $\hat{\rho} = \sum_{ii'} \rho_{ii'} \, |\psi_i\rangle\langle\psi_{i'}|$.

The system states at different energies are transformed independently. Let us consider a system density matrix at a fixed energy subspace of the particle Hilbert space, $\hat{\rho}_E = \sum_{ij} \rho_{ijE} \, |\psi_{i,E}\rangle\langle\psi_{j,E}|$. Then making use of the factorization representation of the evolution operator, see Eq. (A2), one can rewrite the quantum channel in the form,

$$[\Phi(\hat{\rho}_E)]_{jj'} = \sum\nolimits_{ii'} \rho_{ii'E} \, s_{ji,E} s^*_{j'i',E} \, \langle \hat{F}^\dagger_{j'i',E} \hat{F}_{ji,E} \rangle, \qquad \text{(A3)}$$

where the averaging of the reservoir operators $\hat{F}_{ji,E}$ is done with respect to the density matrix $\hat{\pi}$: $\langle \ldots \rangle =$

$\mathrm{tr}\{\hat{\pi}\dots\}$. Now let us consider transformation of $\rho_{ii'E} = \delta_{ii'} = \hat{1}_E$. Then using Eq. (A3) we find:

$$[\Phi(\hat{1}_E)]_{jj'} = \sum\nolimits_i s_{ji,E} s^*_{j'i,E} \langle \hat{F}^\dagger_{j'i,E} \hat{F}_{ji,E} \rangle. \quad \text{(A4)}$$

Averaging Eq. (A1) with respect to the density matrix of the system, moving the left hand side of the averaged equation to the right, and adding the result to the RHS of Eq. (A4), one arrives at Eq. (3) of the text:

$$[\Phi(\hat{1}_E)]_{jj'} = \delta_{jj'} + \sum\nolimits_i s_{ji,E} s^*_{j'i,E} \langle [\hat{F}^\dagger_{j'i,E}, \hat{F}_{ji,E}] \rangle. \quad \text{(A5)}$$

Now we outline the proof of the H-theorem using Eq. (1). To that end we have to calculate $\Phi(\hat{1}) = \Phi(\prod_E \otimes \hat{1}_E)$. Let $[\Phi(\hat{1}_E)]_{jj'} = \delta_{jj'}$ on some subset $\mathcal{E} = \{E\}$. Then using (A4) in the RHS of Eq. (1), we obtain

$$-k_{\mathrm{B}} \sum_{jj',E} \langle \psi_{j,E}|\Phi(\hat{\rho})|\psi_{j',E}\rangle\langle\psi_{j',E}| \log \Phi(\hat{1})|\psi_{j,E}\rangle. \quad \text{(A6)}$$

Accordingly, in Eq. (A6) the terms for which $E \in \mathcal{E}$, become

$$\langle \psi_{j,E}|\Phi(\hat{\rho})|\psi_{j,E}\rangle\langle\psi_{j,E}| \log(1)|\psi_{j,E}\rangle = 0.$$

Now if $\langle\psi_{i,E}|\,\hat{\rho}\,|\psi_{i,E}\rangle \neq 0$ only for $E \in \mathcal{E}$ (therefore, for energies $E \notin \mathcal{E}$, all matrix elements $\langle\psi_{i,E}|\,\hat{\rho}\,|\psi_{j,E}\rangle = 0$), the entire expression (A6) is zero, Eq. (1) yields $\Delta S \geqslant 0$, and the theorem is proved.

Appendix B: Unitarity constraints

Consider a quantum system (particle) which interacts with a quantum reservoir. The evolution of the joint grand system (particle plus reservoir) can be presented in the form,

$$|\psi_i\rangle|n\rangle \to \sum\nolimits_{jm} U_{jm,in}\,|\psi_j\rangle|m\rangle, \quad \text{(B1)}$$

where $|\psi_i\rangle$ and $|n\rangle$ are some orthonormal complete sets of states in the particle and reservoir Hilbert space respectively. The transition amplitudes $U_{jm,in}$ comprise a unitary matrix $\hat{U}$: $\hat{U}^\dagger\hat{U} = \hat{U}\hat{U}^\dagger = \hat{1}$.

We assume no energy exchange between the quantum system and the reservoir. Let us consider a fixed energy subspace E of the particle Hilbert space spanned by a basis $|\psi_{i,E}\rangle$, here index i denotes all remaining degrees of freedom of the particle. Then one can represent the grand evolution operator $\hat{U}$ in the form,

$$\hat{U} = \sum\nolimits_E |\psi_{j,E}\rangle\langle\psi_{i,E}|\, s_{ji,E}\hat{F}_{ji,E}, \quad \text{(B2)}$$

Then the unitarity constraints can be rewritten in the form,

$$\sum\nolimits_i s^*_{ij',E} s_{ij,E} \langle m'|\hat{F}^\dagger_{ij',E}\hat{F}_{ij,E}|m\rangle = \delta_{jj'}\delta_{mm'}, \quad \text{(B3)}$$

$$\sum\nolimits_i s_{ji,E} s^*_{j'i,E} \langle m|\hat{F}_{ji,E}\hat{F}^\dagger_{j'i,E}|m'\rangle = \delta_{jj'}\delta_{mm'}. \quad \text{(B4)}$$

Appendix C: Example of a non-unital channel

Let us inspect a charged particle moving in a three-lead conductor and interacting with the spin via induced magnetic field, see Fig. 1, and, according to the framework of our general consideration, initially disentangled from the spin. In the absence of the external magnetic field the energy exchange is absent and the particle is energy-isolated. The joint scattering states of the particle and spin have the form

$$|\psi^{(\mathrm{in})}_\alpha\rangle|\sigma_0\rangle \to \sum\nolimits_\beta s_{\beta\alpha}\,|\psi^{(\mathrm{out})}_\beta\rangle\,\hat{U}_\beta\hat{U}^\dagger_\alpha|\sigma_0\rangle, \quad \text{(C1)}$$

where $|\psi^{(\mathrm{in/out})}_\alpha\rangle$ is the particle's incoming/outgoing state in the lead α, $s_{\beta\alpha}$ are the components of the scattering matrix of the three lead set up, $|\sigma_0\rangle$ is the initial state of the spin and $\hat{U}_\alpha$ ($\hat{U}^\dagger_\alpha$) is the unitary spin-1/2 rotation of the spin due to outgoing (incoming) electron in the lead α. Then the operators $\hat{F}$ in Eq. (2) are defined as $\hat{F}_{\beta\alpha} = \hat{U}_\beta\hat{U}^\dagger_\alpha$.

We choose spin-1/2 rotations as $\hat{U}_1 = \hat{1}$, $\hat{U}_2 = i\hat{\sigma}_x$ and $\hat{U}_3 = i\hat{\sigma}_y$, where $\hat{\sigma}_x$ and $\hat{\sigma}_y$ are the Pauli matrices. In this case $[\hat{U}_\alpha, \hat{U}_\beta] \neq 0$. Accordingly, $\langle[\hat{F}^\dagger_{\beta'\alpha}, \hat{F}_{\beta\alpha}]\rangle \neq 0$, and the resulting quantum channel is non-unital. Then,

$$\begin{aligned}\Phi(\hat{1}) = \hat{1} &- \big\{|1\rangle\langle 2|\, 2i(s_{11}s^*_{21} + s_{12}s^*_{22})\,\langle\hat{\sigma}_x\rangle + \mathrm{H.c.}\big\} \\ &- \big\{|1\rangle\langle 3|\, 2i(s_{11}s^*_{31} + s_{13}s^*_{33})\,\langle\hat{\sigma}_y\rangle + \mathrm{H.c.}\big\} \\ &+ \big\{|2\rangle\langle 3|\, 2i(s_{22}s^*_{32} + s_{23}s^*_{33})\,\langle\hat{\sigma}_z\rangle + \mathrm{H.c.}\big\}. \quad \text{(C2)}\end{aligned}$$

To simplify this equation, we assume time-reversal symmetry of the electron scattering, $s_{ij} = s_{ji}$, and exploiting the following representation[16] of the elastic scattering matrix,

$$s_{ii} = \frac{1}{2} s^*_{jk} s_{ij} s_{ik} \Big(\frac{1}{T_{jk}} - \frac{1}{T_{ij}} - \frac{1}{T_{ik}}\Big), \quad i \neq j \neq k, \quad \text{(C3)}$$

where $T_{ij} = |s_{ij}|^2$ are transmission probabilities, we obtain Eq. (5).

Appendix D: One-dimensional scattering

From the unitarity of the elastic scattering matrix it follows $|s_{\mathrm{RR}}|^2 + |s_{\mathrm{LR}}|^2 = 1$, $|s_{\mathrm{RL}}|^2 + |s_{\mathrm{LL}}|^2 = 1$, $s^*_{\mathrm{LL}}s_{\mathrm{LR}} + s^*_{\mathrm{RL}}s_{\mathrm{RR}} = 0$, and $s_{\mathrm{RL}}s^*_{\mathrm{LL}} + s^*_{\mathrm{LR}}s_{\mathrm{RR}} = 0$. The unitarity of the joint particle-reservoir gives us

$$s^*_{\mathrm{LL}}s_{\mathrm{LR}}\,\langle n''|n\rangle + s^*_{\mathrm{RL}}s_{\mathrm{RR}}\,\langle n|n'\rangle = 0. \quad \text{(D1)}$$

Hence $\langle n''|n\rangle = \langle n|n'\rangle$ and

$$s^*_{\mathrm{LL}}s_{\mathrm{RL}}\langle n''|n\rangle + s^*_{\mathrm{LR}}s_{\mathrm{RR}}\langle n|n'\rangle = 0. \quad \text{(D2)}$$

Let us consider the transformation of the identity operator $\hat{1} = |\mathrm{R}\rangle\langle\mathrm{R}| + |\mathrm{L}\rangle\langle\mathrm{L}|$ if reservoir initially rests in a

state $|n\rangle\langle n|$,

$$\hat{1} \to |\mathrm{R}\rangle\langle\mathrm{R}|\,(|s_{\mathrm{RL}}|^2 + |s_{\mathrm{RR}}|^2) + |\mathrm{L}\rangle\langle\mathrm{L}|\,(|s_{\mathrm{LL}}|^2 + |s_{\mathrm{LR}}|^2) + \big\{|\mathrm{R}\rangle\langle\mathrm{L}|\,(s^*_{\mathrm{LL}}s_{\mathrm{RL}}\langle n''|n\rangle + s^*_{\mathrm{LR}}s_{\mathrm{RR}}\langle n|n'\rangle) + \mathrm{H.c.}\big\}. \quad \text{(D3)}$$

One can see that the off-diagonal terms $|\mathrm{R}\rangle\langle\mathrm{L}|$ of the transformed identity operator vanish due to unitarity constraint, see Eq. (D2), while the diagonal elements sum to the identity operator and hence $\Phi(\hat{1}) = \hat{1}$. The same arguments are valid if initially reservoir rests in an arbitrary mixed state non-entangled with the particle.

The same arguments can be given in a more general form, exploiting the H-theorem on page 2. Making use of the scattering states [Eqs. (6)–(7)] one can construct the joint evolution operator at a given energy of the incoming particle,

$$\hat{U} = s_{\mathrm{LR}}|\mathrm{L}\rangle\langle\mathrm{L}| \otimes \hat{1} + s_{\mathrm{RL}}|\mathrm{R}\rangle\langle\mathrm{R}| \otimes \hat{1} + s_{\mathrm{RR}}|\mathrm{R}\rangle\langle\mathrm{L}| \otimes \hat{F}_{\mathrm{RR}} + s_{\mathrm{LL}}|\mathrm{L}\rangle\langle\mathrm{R}| \otimes \hat{F}_{\mathrm{LL}},$$

which transforms the incoming particle states into outgoing states. Here, first two terms corresponds to the forward scattering of the particle which does not change the reservoir state. The last two terms accounts for the backscattering events which do change the reservoir state. The joint unitarity constraint $\hat{U}^\dagger\hat{U} = \hat{1}$ gives,

$$(|s_{\mathrm{LR}}|^2\hat{1} + |s_{\mathrm{RR}}|^2\hat{F}^\dagger_{\mathrm{RR}}\hat{F}_{\mathrm{RR}})|\mathrm{L}\rangle\langle\mathrm{L}| + (|s_{\mathrm{RL}}|^2\hat{1} + |s_{\mathrm{LL}}|^2\hat{F}^\dagger_{\mathrm{LL}}\hat{F}_{\mathrm{RR}})|\mathrm{R}\rangle\langle\mathrm{R}| + \big\{\big(s^*_{\mathrm{LL}}s_{\mathrm{LR}}\hat{F}^\dagger_{\mathrm{LL}} + s^*_{\mathrm{RL}}s_{\mathrm{RR}}\hat{F}_{\mathrm{RR}}\big)|\mathrm{R}\rangle\langle\mathrm{L}| + \mathrm{H.c.}\big\} = \hat{1}. \quad \text{(D4)}$$

The unitarity of the elastic scattering amplitudes $s^*_{\mathrm{RL}}s_{\mathrm{RR}} + s^*_{\mathrm{LL}}s_{\mathrm{LR}} = 0$ vanish the off-diagonal terms $|\mathrm{L}\rangle\langle\mathrm{R}|$ and $|\mathrm{R}\rangle\langle\mathrm{L}|$ on the Eq. (D4) if

$$\hat{F}_{\mathrm{RR}} = \hat{F}^\dagger_{\mathrm{LL}}, \quad \text{(D5)}$$

while the normalization condition $|s_{\mathrm{LR}}|^2 + |s_{\mathrm{RR}}|^2 = 1$ and $|s_{\mathrm{RL}}|^2 + |s_{\mathrm{LL}}|^2 = 1$ gives,

$$\hat{F}^\dagger_{\mathrm{RR}}\hat{F}_{\mathrm{RR}} = \hat{1}, \quad \hat{F}^\dagger_{\mathrm{LL}}\hat{F}_{\mathrm{LL}} = \hat{1}, \quad \text{(D6)}$$

and hence $\hat{F}_{\mathrm{RR}}$ and $\hat{F}_{\mathrm{LL}} = \hat{F}^\dagger_{\mathrm{RR}}$ are unitary operators. It follows then that $[\hat{F}_{\beta'\alpha}, \hat{F}_{\beta\alpha}] = 0$ for $\alpha,\ \beta,\ \beta' \in \{\mathrm{L}, \mathrm{R}\}$ and the scattering process [Eqs. (6)–(7)] describes an unital quantum channel.

[1] L. Boltzmann, *Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen*, Wiener Berichte **75**, 62–100 (1872).

[2] L. Boltzmann, *Entgegnung auf die wärme-theoretischen Betrachtungen des Hrn. E. Zermelo*, Annalen der Physik (Leipzig) **57**, 773–784 (1896) [Translated and reprinted in S. G. Brush, *Kinetic Theory 2* (Pergamon Elmsford, 1966)].

[3] J. von Neuman, *Beweis des Ergodensatzes und des H-Theorems in der neuen Mechanik*, Zeitschrift für Physik **57**, 30–70 (1929) [J. Von Neumann, *Proof of the ergodic theorem and the H-theorem in quantum mechanics*, Eur. Phys. J. H **35**, 201–237 (2010)].

[4] J. Gemmer, M. Michel, and G. Mahler, *Quantum thermodynamics*, (Springer, 2009).

[5] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information*, (Cambridge Univ. Press, 2011).

[6] A. S. Holevo, *Quantum systems, channels, information. A mathematical introduction*, (De Gryter, 2012).

[7] A. S. Holevo, *The entropy gain of infinite-dimensional quantum channels*, Doklady Math. **82**, 730–731 (2010).

[8] A. Wehrl, *General properties of entropy*, Rev. Mod. Phys. **50**, 221–260 (1978).

[9] G. G. Amosov, *Estimating the output entropy of a tensor product of two quantum channels*, Theor. Math. Phys. **182**, 397–406 (2015).

[10] G. Argentieri, F. Benatti, R. Floreanini, and M. Pezzutto, *Violations of the second law of thermodynamics by a non completely positive dynamics*, Europhys. Lett. **107**, 50007 (2014).

[11] G. B. Lesovik, *On the law of increasing entropy and the cause of the dynamics irreversibility of quantum systems*, JETP Lett. **98**, 184–189 (2013).

[12] Y. Aharonov, L. Davidovich, and N. Zagury, *Quantum random walks*, Phys. Rev. A **48**, 1687 (1993).

[13] A. M. Childs, E. Farhi, and S. Gutmann, *An example of the difference between quantum and classical random walks*, Quant. Inf. Proc. **1**, 35 (2002).

[14] V. Coffman, J. Kundu, and W. K. Wootters, *Distributed entanglement*, Phys. Rev. A **61**, 052306 (2000).

[15] L. P. Pitaevskii and E. M. Lifshitz, *Physical Kinetics*, (Butterworth-Heinenann, 1981).

[16] G. B. Lesovik, T. Martin, and G. Blatter, *Electronic entanglement in the vicinity of a superconductor*, Euro. Phys. B **24**, 287–290 (2001).